




SUB-LABEL B:

For Gardening / Residential Use





ACTIVE INGREDIENTS:

Pyrethrins	1.40%
Azadirachtin.....	1.20%
OTHER INGREDIENTS.....	97.40%
TOTAL	100.00%

Contains 0.10 lbs of azadirachtin and 0.11 lbs of pyrethrins per gallon.

**KEEP OUT OF REACH OF CHILDREN
CAUTION PRECAUCIÓN**

**Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail).**

FIRST AID	
IF IN EYES	<ul style="list-style-type: none">• Hold eye open and rinse slowly and gently with water for 15-20 minutes.• Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.• Call a poison control center or doctor for treatment advice.
IF ON SKIN OR CLOTHING	<ul style="list-style-type: none">• Take off contaminated clothing.• Rinse skin immediately with plenty of water for 15-20 minutes.• Call a poison control center or doctor for treatment advice.
IF SWALLOWED	<ul style="list-style-type: none">• Call poison control center or doctor immediately for treatment advice.• Have person sip a glass of water if able to swallow.• Do not induce vomiting unless told to do so by the poison control center or doctor• Do not give anything by mouth to an unconscious person.
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information regarding medical emergencies or pesticide incidents, call 1-888-740-8712.	

LAM INTERNATIONAL CORPORATION

117 S. Parkmont; P.O. Box 4109-Butte, MT 59702; Ph: (406)782-2386; Fax:(406)782-9912

EPA Reg. No. 82074-XX EPA Est. Number 65626-MT-02

Net Contents: _____

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash hands thoroughly with plenty of soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Wear protective eyewear. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals. Wear long-sleeved shirt and long pants, socks, shoes and chemical-resistant gloves (such as barrier laminate, nitrile rubber, neoprene rubber, or viton).

ENVIRONMENTAL HAZARDS

To protect the environment, do not allow pesticide to enter or run off into storm drains, drainage ditches, gutters, or surface waters. Applying this product in calm weather when rain is not predicted for the next 24 hours will help to ensure that wind or rain does not blow or wash pesticide off the treatment area.

This product is toxic to aquatic organisms, including fish and invertebrates. Drift and run-off may be hazardous to aquatic organisms in water adjacent to treated areas. This product may contaminate water through run-off. This product has a potential for run-off for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce run-off that contains this product.

This product is highly toxic to bees exposed to direct treatment on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds while bees are foraging the treatment area.

Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash-waters or rinsate. See Directions for Use for additional precautions and restrictions.

PHYSICAL OR CHEMICAL HAZARDS

Combustible: Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling

RESTRICTIONS:

- Apply this product only as specified on this label.
- Do not contaminate food or feedstuffs.
- Do not apply this product in a way that will contact workers or other persons, either directly or through spray drift.
- Do not allow adults, children, or pets to enter the treated area until sprays have dried.
- Do not remain in treated area. Exit area immediately and remain outside the treated area until vapors, mists and aerosols have dispersed.
- Do not wet plants to the point of runoff or drip.
- Do not apply directly to or near water, storm drains or drainage ditches. Do not apply when windy.
- Rinse applicator over lawn or garden area only.
- To avoid possible harm to honey bees, it is advisable to apply in the early morning or late evening hours.
- Do not apply more than 1 time per day.
- Do not apply more than 10 times per season.
- In case of extreme pest pressure, do not reapply within 24 hours.
- Do not reapply within 3 days except under extreme pest pressure.
- Do not make applications during rain.
- Not for use in indoor or outdoor residential misting systems.

DIRECTIONS

Phytotoxicity: SPHYNX has been evaluated for phytotoxicity on a wide range of plants. However, since testing on all varieties of all plants is not feasible, nor is testing of all possible combinations or sequences of pesticide sprays including fertilizers, surfactants and adjuvants, before making tank mix combinations with SPHYNX, or before making widespread applications, it is recommended to treat a limited number of plants and observe for phytotoxicity over a 10-day period. It is further recommended that spray equipment used to apply MGK - 2905 be thoroughly cleaned before use.

Mode of Action: SPHYNX kills target pests quickly by contact or ingestion. It also kills listed insects by interfering with the molting process and as a adulticide. It is effective on all larva stages, pupae, and adults.

Honey Bees: To avoid possible harm to honey bees, it is advisable to apply in early morning or late evening hours.

RATE CHART:

		<i>Most commonly used rate</i>	<i>Rates for treating high populations of adults and/or hard to kill insects</i>	
SPHYNX	1 Pint per acre 16 fl. oz. (473 ml)	2 Pints per acre 32 fl. oz. (946 ml)	3 Pints per acre 48 fl. oz. (1.42 L)	3.5 Pints per acre 56 fl. oz. (1.66 L)
Acres per Quart	2	1	0.67	0.57
Acres per Gallon	8	4	2.7	2.3

DILUTION RATES

CONVENTIONAL EQUIPMENT: In sufficient water for thorough coverage. Dilution in a minimum of 30 gallons (114 L) of water per acre is recommended

HAND SPRAYER: 1-2 fluid ounces (30-60 ml) of SPHYNX per gallon (3.8 L) of water

GREENHOUSE: Dilute 53-107 fl. oz. (1567-3164 ml) with 100 gallons (378.54 L) of water for applications with conventional hydraulic sprayers, or 1 to 2 fl. oz. (30-60 ml) per one gallon (3.8 L) of water, or applications with compressed sprayers. Use 2.3 gallons (8.71 L) of spray solution per 1,000 square feet (93 m²).

MIXING DIRECTIONS

USED ALONE:

- Mix only enough for immediate use.
- Shake SPHYNX well before using.
- Dilute SPHYNX in sufficient water to obtain thorough coverage.
- Fill clean spray tank 1/2 to 3/4 of the water to be sprayed and begin agitation.
- Add the appropriate amount of SPHYNX to the spray tank.
- Fill the tank with the remaining water and agitate thoroughly.
- Apply product promptly after mixing.
- Complete coverage of all leaf surfaces is essential for optimum results.
- If the mixture is not applied immediately after mixing, agitate before application.

APPLICATION DIRECTIONS:

Spraying should begin when listed insects first appear. Do not wait until plants are heavily infested. Repeat application as required to maintain effective kill, but not more than every 5-7 days. For foliar application, apply SPHYNX in sufficient spray volume and with adequate spray pressure to ensure complete and thorough coverage of all plant surfaces including both the top and bottom of leaves. Do not wet plants to the point of runoff or drip. Do not apply when wind speed favors drift beyond the area intended for treatment. When pest pressure is extreme or plant canopy is dense, use higher rates and do not reapply within 24 hours. If possible, apply in the early morning or evening hours. The reduced UV exposure and lower temperatures will increase the performance and reduce the impact on pollinators.

INSECTS CONTROLLED

SPHYNX is labeled for the control of insects including:

Aphids including: Apple Aphids, Alfalfa Aphids, Artichoke Aphids, Bean Aphids, Black Maringed Aphids, Black Bean Aphids, Black Peach Aphids, Blue Alfalfa Aphids, Cabbage Aphids, Cotton / Melon Aphids, Cowpea Aphids, European Asparagus Aphids, Filbert Aphids, Foxglove Aphids, Green Peach Aphids, Lettuce Aphids, Lettuce Root Aphids, Melon Aphids, Pea Aphids, Potato Aphids, Rose Aphids, Spotted Alfalfa Aphids, Willow Carrot Aphids

Armyworms, Caterpillars and Loopers including: Alfalfa Caterpillars Artichoke Plume Moths, Bagworms, Beet Armyworms, Black Cutworms, Budworms, Cabbage Loopers, Cankerworms, Carpenterworms, Citrus Cutworms, Corn Earworms, Cross-striped cabbageworms, Cutworms, Diamondback moths, Eastern Tent Caterpillars, Fall Armyworms, Fall Cankerworms, Fall Webworms, Filbert Worms, Fireworms, Forest Tent Caterpillars, Garden Webworms, Grapefruit Worms, Grape Leaf Skeletonizers, Green Fruit Worms, Hickory Shuckworms, Hornworms, Imported Cabbageworms, Lawn Armyworms, Lesser Webworm, Loopers, Melonworms, Navel Orangeworms, Oriental Fruit Moths, Pecan Nut Case bearers, Rindworms, Sod Webworms, Southern Armyworms, Soybean Loopers, Saltmarsh Caterpillars, Tent Caterpillars, Tobacco Budworms, Tomato Hornworms, Tomato Fruitworms, Tomato Pinworms, Yellow striped, Armyworms, Walnut Caterpillars Webworms, Western Yellow-Striped Armyworms, Western Grapeleaf Skeletonizers.

Beetles and Weevils including: Alfalfa Weevils, Asparagus Beetles, Bean Beetles, Bean Leaf Beetles, Black Vine Weevils, Blister Beetles, Boll Weevils, Carrot Weevils, Chestnut Weevils Clover Weevils Colorado Potato Beetles, 12-Spotted Cucumber Beetles, Cucumber Beetles, Darkling Beetles (lesser meal worms), Egyptian Alfalfa Weevils, Elm Leaf Beetles, Flea Beetles, Fuller Rose Beetles, Grape Bud Beetles, Japanese Beetles, June Beetles, Mexican Bean Beetles, Navel Orangeworms (NOW), Pecan Weevils, Pink Bollworms, Potato Flea Beetles, Rice Weevils, Rose Chafers, Saw-Toothed Grain Beetles, Strawberry Beetles, Twig Girdlers, All other beetles and weevils.

Leafrollers including: Blueberry Leafrollers, Filbert Leafrollers, Fruit Tree Leafrollers, Grape Leafrollers, Oblique Banded Leafrollers, Omnivorous Leafrollers, Orange Tortrix, Western Avocado Leafrollers.

Borers such as: European Corn Borers, Pacific Flatheaded Borers, Peach Tree Borers, Peach Twig Borers, Squash Vine Borers, Shotbole Borers, Branch and Twig Borers.

Flies: Australian Sod Flies, Caribbean Fruit Flies, Crane Flies, Fruit Flies, Fungus Gnats, Hessian Flies, Mediterranean Fruit Flies, Melon Flies, Mushroom Flies, Oriental Fruit Flies, Olive Fruit Flies, Sawflies, Shore Flies, Vinegar Flies, Walnut Husk Flies.

Leafhoppers & Sharpshooters: Aster Leafhoppers, Beet Leafhoppers, Glassy-winged Sharpshooters, Grape Leafhoppers, Potato Leafhoppers, Variegated Leafhoppers, Three-Cornered Alfalfa hoppers.

Leafminers: Citrus Leafminers, Holly Leafminers, Serpentine Leafminers, Vegetable Leafminers.

Midges (plant pests): Millipedes, Onion Maggots, Plant Bugs, Proba Bugs, Scale Insects, Silverfish, Skippers, Soft Scales, Spider Mites, Sowbugs, Spiders (except Black Widow and Brown Recluse Spiders), Springtails, Squash Bugs, Stink Bugs, Tarnished Plant Bugs, Spittle Bugs, Wire worms, European Chafers, Northern Masked Chafers, Southern Masked Chafers, Western Boxelder Bugs.

Moths: Artichoke Plume Moths, Codling Moths, Diamondback Moths, European Pine Tip Moths, Grape Berry Moths, Gypsy Moths (adult & larvae), Indian Meal Moths, Mediterranean Flour Moths, Pine Tip Moths, Tussock Moths

Whiteflies: Greenhouse Whiteflies, Silverleaf Whiteflies, Sweetpotato Whiteflies,

Other: Ants (except Pharaoh, Harvester, Carpenter and Fire Ants), Apple Maggots, Billbugs, Brown Marmorated Stinkbugs, Cabbage Maggots, Clover Mites, Cutworms, Crickets, Dichondra Flea Beetles, Earwigs, Firebrats, False Chinch Bugs, Garden Symphytan, Garden Tortrix, Glassy Winged Sharpshooters, Grasshoppers, Harlequin Bugs, Grape Phylloxera, Katydid, Leaf-footed Plant Bug, Lace Bugs, Leaf tiers, Lice, Lygus, Lace bug, Mealybugs (all)

Psyllids: Pear Psylla

Thrips: Avocado thrips, Citrus Thrips, Flower Thrips, Greenhouse Thrips, Thrips Palmi, Northern Masked Chafers, Southern Masked Chafers, Western Flower Thrips

FOR USE ON VEGETABLES, FRUITS, HERBS, AND SPICES IN RESIDENTIAL GARDENS:

ROOT AND TUBER VEGETABLES: Including: Arracacha, Arrowroot, Purple Arrowroot, Japanese Artichokes, Jerusalem Artichokes, Garden Beets, Sugar Beets, Edible Burdock, Edible Canna, Carrots, Cassava (bitter or sweet), Celeriac, Celery Root, Chayote (root), Chervil (turnip rooted), Chicory, Chufa, Dasheen (Taro), Ginger, Ginseng, Horseradish, Jicama, Leren, Parsley (turnip rooted), Parsnips, Potatoes, Radishes, Japanese Radishes (Daikon), Rutabaga, Salsify (oyster plant, black, Spanish), Skirret, Sweet Potatoes, Tanier, Turmeric, Turnips, Yam Beans (jicama, manioc pea), Yams (true).

LEAVES OF ROOT AND TUBER VEGETABLES: Including: Garden Beets, Sugar Beets, Edible Burdock, Carrots, Cassava (bitter and sweet), Celeriac (celery root), Chervil (turnip-rooted), Chicory, Dasheen (Taro), Parsnips, Radishes, Oriental Radishes (Daikon), Rutabaga, Salsify (black), Sweet Potatoes, Tanier, Turnips, Yams (true).

BULB VEGETABLES: Including: Garlic, Great-headed Garlic, Leeks, Onions (bulb and green), Shallots, Welch.

LEAFY VEGETABLES: Including: Amaranth (Leafy Amaranth, Chinese Spinach, Tampala), Arugula, Cardoon, Celery, Chinese Celery, Celtuce, Chervil, Corn Salad, Chrysanthemum (edible-leaved), Chrysanthemum (garland), Cress (garden, water), Upland Cress (yellow rocket, winter cress), Dandelion, Dock (sorrel), Endive (escarole), Fennel (Florence), Leeks, Lettuce (head and leafy), Mustard Greens, Orach, Parsley, Purslane (garden & winter), Radicchio, Rhubarb, Spinach, Fine Spinach (Malabar, Ceylon), Spinach (New Zealand), Swiss Chard, Turnip Greens, Watercress.

BRASSICA (COLE) LEAFY VEGETABLES: Including: Broccoli, Chinese Broccoli (Gai Lan), Broccoli raab (Rapini), Brussels Sprouts, Cabbage, Chinese Cabbage (Bok Choy), Chinese Cabbage (Napa), Chinese Mustard Cabbage (Gai Choy), Cauliflower, Cavalo broccolo, Collards, Kale (Flowering, Chinese), Kohlrabi, Mizuna, Mustard Greens, Mustard Spinach, Rape Greens.

LEGUME VEGETABLES (SUCCULENT OR DRIED): Including: Adzuki Beans, Field Beans, Kidney Beans, Lima Beans, Moth Beans, Mung Beans, Navy Beans, Pinto Beans, Rice Beans, Runner Beans, Snap Beans, Tepary Beans, Urd Beans, Wax Beans, Asparagus Beans, Black-eyed Peas, Catjang, Chinese Longbeans, Cowpeas, Crowder Peas, Southern Peas, Yard-Longbeans, Broad Beans, (Fava Beans), Chick Peas (Garbanzo Beans), Guar, Jackbean (Sword Bean), Lablab Bean (Hyacinth Bean), Lentils, Peas (Garden Peas, Field Peas, Sugar Snap Peas, English Pea, Snow Pea), Pigeon Peas, Soybeans, Sweet Lupin Beans, White Lupin Beans, White Sweet Lupin, Sword Bean.

FOLIAGE OF LEGUME VEGETABLES: Including: Plant part of any legume vegetable included in the legume vegetable group that will be used as animal feed including any variety of Beans, Field Peas, Soybeans.

FRUITING VEGETABLES: Including: Eggplant, Ground Cherry, Okra, Pepinos, Pepper (Bell Pepper, Chili Pepper, Cooking Peppers, Pimentos, Sweet Peppers), Tomatillo, Tomatoes.

CUCURBIT VEGETABLES: Including: Balsam Apple, Balsam Pear (Bitter Melon), Chayote (fruit), Chinese Waxgourd (Chinese preserving melon), Chinese Cucumber, Citron Melon, Cucumber, Gherkin, Edible Gourds, Mangoes, Melons (including hybrids, Cantaloupe, Casaba, Charentais, Crenshaw, Golden Pershaw Melon, Honeydew Melons, Honey Balls, Mango Melon, Muskmelon, Persian Melon, Pineapple Melon, Santa Claus Melon, Snake Melon, Pumpkin, Squash (summer & winter), Watermelon (OP & hybrids, seeded and seedless).

CITRUS FRUITS: Including: Calamondin, Citrus Citron, Citrus Hybrids, Grapefruit, Kumquats, Lemons, Limes, Mandarin (Tangerine), Orange (sweet & sour), Pummelo, Satsuma Mandarin, (Citrus spp. Includes Chironja, Tangelos, Tangors), Unique Fruit, White Sapote.

POME FRUITS: Apple, Crabapple, Loquat, Mayhaw, Oriental Pear, Pear, Quince.

STONE FRUITS: Apricot, Cherry (sweet & sour), Nectarine, Peach, Plum, Prune, Chickasaw Plum, Damson Plum, Japanese Plum, Plumcot.

SMALL FRUITS AND BERRIES: Blackberry, Blueberry, Boysenberries, Cranberry, Currant, Dewberry, Elderberry, Gooseberry, Guava, Grape, Huckleberry, Loganberry, Olallie Berry, Raspberry (black & red), Strawberry, Youngberry.

TREE NUTS: Almond, Beech Nut, Brazil Nut, Butternut, Cashew, Chestnut, Chinquapin, Filbert (hazelnut), Hickory nut, Macadamia Nut (Bush Nut), Pecan, Pistachio, Walnut (Black, English, Persian).

TROPICAL FRUITS: Papaya, Black Sapote, Canistel, Mango, Sapodilla, Star Apple, Guava, Biriba.

CEREAL GRAINS: Barley, Buckwheat, Corn (sweet and field), Millet, Proso, Oats, Pearl Millet, Popcorn, Rice, Rye, Sorghum (Milo), Teosine, Triticale, Wheat, Wild Rice.

FORAGE, FODDER AND STRAW OF CEREAL GRAINS: Barley; buckwheat; corn (sweet and field); millet; proso; oats; pearl; popcorn; rice; rye; sorghum (milo); teosine; triticale; wheat; wild rice.

GRASSES FOR SEED, FORAGE, FODDER AND HAY: any grass (all Gramineacea), (green or cured), except sugarcane and those listed in the cereal grains group) that will be fed to or grazed by livestock, all Pasture and Range Grasses and Grasses grown for hay and silage, Sudangrass, Bermuda Grass, Bluegrass, Bromegrass, Fescue, Orchard, Timothy.

NON-GRASS ANIMAL FEEDS: Alfalfa, Velvet Bean, Clover (White, Ladino, Red), Kudzu, Lespedeza, Lupine, Sainfoin, Trefoil, Crown Vetch, Milk Vetch.

HERBS AND SPICES: Allspice, Angelica, Anise (anise seed), Anise [star], Annatto (seed), Balm (lemon balm), Basil, Borage, Burnet, Chamomile, Caper buds, Caraway, Caraway [black], Cardamom, Cassia bark, Cassia buds, Catnip, Celery seed, Chervil (dried), Chicory, Chive, Chive [Chinese], Cinnamon, Clary, Clove buds, Coriander (cilantro or Chinese parsley) (leaf), Coriander (cilantro) (seed), Costmary, Cilantro (leaf), Cilantro (seed), Cumin, Curry (leaf), Dandelion, Dill (dillweed), Dill (seed), Fennel (common), Fennel [Florence] (seed), Fenugreek, Grains of paradise, Horehound, Hyssop, Juniper berry, Lavender, Lemongrass, Lovage (leaf), Lovage (seed), Mace, Marigold, Marjoram (includes sweet or annual marjoram, wild marjoram or oregano, and pot marjoram), Mint, Mustard (seed), Nasturtium, Nutmeg, Oregano, Mint, Paprika, Parsley (dried), Pennyroyal, Pepper [black], Pepper [white], Peppermint, Poppy (seed), Rosemary, Rue, Saffron, Sage, Savory [summer and winter], Sweet bay (bay leaf), Tansy, Tarragon, Thyme, Vanilla, Wintergreen, Woodruff, Wormwood.

OIL SEED GROUP: Cottonseed, Jojoba.

ADDITIONAL CROPS: Acerola, Artichoke, Asparagus, Avocado, Atemoya, Bananas, Barbados Cherry, Birdseed, Cacao, Canistel, Carob, Cherimoya, Cocoa, Coffee, Custard Apple, Dates, Durian (Jackfruit), Edible Flowers, Feijoa, Figs, Globe Artichoke, Guayule, Hops, Jaboticaba, Llama, Longan, Lychee, Mangoes, Mamey Sapote, Mushroom, Okra, Olives, Palm, Papaya, Passion fruit, Peanuts, Persimmon, Pineapple, Pomegranate, Pulasan, Rambutan, Safflowers, Sapodilla, Sesame, Soursop and Biriba, Spanish Lime, Star Apple, Sugar Cane, Star Fruit, Sugar Apple, Sunflower, Tamarillo, Tea, Tobacco, Water chestnut and Wax Jambu.

FOR USE ON GROWING PLANTS (OUTDOORS):

BEDDING PLANTS, FLOWERS, POTTED PLANTS AND FOLIAGE: Actinopteris, Aglaonema, Allamanda, Algerian Ivy, Alocasia, Anthurium, Aphelandra, Artemisia, Aster, Aucuba Illex, Azalea, Baby's Breath, Begonia, Bougainvillea, Boston Fern, Boxwood, Brachycome, Cacti, Calabrese, Caladium, Calla, Calathea, Calendula, Carnation, Chrysanthemum, Coleus, Columbine, Dahlia, Daisy, Daylily, Delphinium, Dianthus, Dieffenbachia, Dusty Miller, Easter Lily, English Ivy, Euphorbia, Fern, Ficus, Foxglove, Freesia, Fuchsia, Gaillardia, Gardenia, Geranium, Gerbera, Gladioli, Gypsophila, Hedera, Hibiscus, Impatiens, Iris, Lily, Mandevilla, Marigold, Nasturtium, Pansy, Pelargonium, Peony, Peperomia, Petunia, Philodendron, Phlox, Photinia, Pittosporum, Pinks, Poinsettia, Pothos, Portulaca, Rosemary, Rose, Rubber plant, Salvia, Schefflera, Sedum, Sempervivum, Snapdragon, Spathiphyllum, Stock, Syngonium, Verbena, Vinca, Wander Jew, Zinnia.

ORNAMENTALS: African Violet, Ageratum, Arborvitae, Aster, Aucuba Illex, Azalea, Begonia, Boxwood, Cacti, Calceolaria, Calendula, Calla, Camellia, Carnation, Ceanothus, Chrysanthemum, Cineraria, Coleus, Cotoneaster, Cyclamen, Cypress, Daffodil, Dahlia, Delphinium, Dogwood, Elm, Eucalyptus, Fern, Ficus, Foliage Plants, Fuchsia, Gardenia, Geranium, Gladiolus, Gloxinia, Gypsophila, Holly, Hyacinth, Hydrangea, Iris, Ivy, Lily, Maidenhair Fern, Marigold, Narcissus, Orchid, Palm, Pansy, Peony, Pelargonium, Petunia, Philodendron, Phlox, Photinia, Pine, Pyracantha, Rhododendron, Roses, Rubber Plant, Snapdragon, Stock, Sweet Pea, Tulips, Viburnum, Wandering Jew, White Cedar, White Pine, Yew, Yucca, Zinnia

TREES AND SHRUBS: Andromeda, Arborvitae, Ash, Austrian Pine, Azalea, Beech, Birch, Birds Nest Spruce, Blue Spruce, Boxwood, Butternut, Cedar, Chamaecyparis, Cherry, Crabapple, Cotoneaster, Cyprus, Dogwood, Douglas Fir, Elm, Euonymus, Fir, Firethorn, Forsythia, Hackberry, Hawthorn, Hemlock, Hickory, Holly, Honey Locust, Horse Chestnut, Juniper, Larch, Laurel, Lilac, Linden, London Plane, Magnolia, Mandevilla, Maple, Mimosa, Mountain Ash, Myrtle, Oak, Pachysandra, Peach, Pine, Planetree, Poplar, Privet, Quince, Spruce, Sycamore, Tulip Tree, Viburnum, Willow.

TURF GRASS: Residential Yards and other Recreational Turf Areas.

STORAGE AND DISPOSAL:

Do not contaminate water, food, or feed by storage or disposal.

PESTICIDE STORAGE: Store in a cool, dry place away from heat or open flame in an area that is inaccessible to children and animals.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste facility.

CONTAINER HANDLING: Nonrefillable container. Do not reuse or refill this container.

If empty: Offer for recycling, if available.

If partially filled: Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

CONDITIONS OF SALE

This product conforms to the description set forth on this label and is reasonably fit for the purposes described herein when used according to the label directions and specified conditions. The manufacturer disclaims any and all other express or implied warranties of merchantability and fitness for particular purpose. To the extent consistent with applicable law, buyers and users shall assume all risk and responsibility for potential loss or damage if this product is used, stored, handled or applied in a manner inconsistent with this labeling. To the extent permitted by law, manufacturer shall not be liable for more than the purchase price for the quantity involved including incidental, consequential or special damages.